

References

1. Liu et al. Association of a functional cytochrome P450 4F2 haplotype with urinary 20-HETE and hypertension. *J. Am. Soc. Nephrol.* 19, 714-721, 2008.
2. Meseguer et al. Kidney androgen-regulated protein transgenic mice show hypertension and renal alterations mediated by oxidative stress. *Circulation.* 119, 1908-1917, 2009.
3. Dolegowska, B., Blogowski, W., Domanski L. Is it possible to predict the early post-transplant allograft function using 20-HETE measurements? A preliminary report. *Transpl Int.* 22, 546-553, 2009.
4. Liu et al. Overexpression of cytochrome P450 4F2 in mice increases 20-hydroxyeicosatetraenoic acid production and arterial blood pressure. *Kidney. International.* 75, 1288-1296, 2009.
5. Wang et al. Selective inhibitors of CYP2J2 related to terfenadine exhibit activity strongly against human cancers *in vitro* and *in vivo*. *J. Pharmacol. Exp. Ther.* 329, 908-918, 2009.
6. Cuez, Malik, Tunctan et al. A synthetic analogue of 20-HETE, 5,14-HEDGE, reverses endotoxin-induced hypotension via increased 20-HETE levels associated with decreased iNOS protein expression and vasodilator prostanoid production in rats. *Basic Clin. Pharmacol. Toxicol.* 106, 378-388, 2010.
7. Malik et al. 2,3',4,5'-Tetramethoxystilbene prevents deoxycorticosterone-salt-induced hypertension: contribution of cytochrome P-450 1B1. *Am. J. Physiol. Heart Circ. Physiol.* 299, H1891-H1901, 2010.
8. Malik et al. Cytochrome P450 1B1 contributes to angiotensin II-induced hypertension and associated pathophysiology. *Hypertension* 56, 667, 2010.
9. Tunctan et al. Contribution of vasoactive eicosanoids and nitric oxide production to the effect of selective cyclooxygenase-2 inhibitor, NS-398, on endotoxin-induced hypotension in rats. *Basic Clin. Pharmacol. Toxicol.* 107, 877-882, 2010.
10. Imaizumi et al. L-4F differentially alters plasma levels of oxidized fatty acids resulting in more anti-inflammatory HDL in mice. *Drug Metab. Letters*, 4, 139-148, 2010.
11. Cervenka, Kramer, Falck, Imig, Hammock et al. Combined inhibition of 20-HETE formation and of EET degradation attenuates hypertension and hypertension-induced end-organ damage in Ren-2 transgenic rats. *Clinical Science* 118, 617-632. 2010.
12. Cervenka, Kramer, Falck, Imig et al. Intrarenal CYP-450 metabolites of arachidonic acid in the regulation of the nonclipped kidney function in two-kidney, one-clip Goldblatt hypertensive rats. *J Hypertens* 28, 582-593, 2010.
13. Hu, Wang et al. Peripheral and central augmentation indexes in relation to the CYP4F2 polymorphisms in Chinese. *J Hypertens* 29, 501-508, 2011.
14. Buharalioglu et al. Piroxicam reverses endotoxin-induced hypotension in rats: contribution of vasoactive eicosanoids and nitric oxide. *Basic Clin Pharmacol Toxicol* 186–194, 2011.
15. Na-Bangchang et al. Study on the association between environmental cadmium exposure, cytochrome P450-mediated 20-HETE, heme-oxygenase-1 polymorphism and hypertension in Thai population residing in a malaria endemic areas with cadmium pollution. *Environ Toxicol Pharmacol* 31, 416-426, 2011.
16. Grande et al. Increased oxidative stress, the renin-angiotensin system, and sympathetic overactivation induce hypertension in kidney androgen-regulated protein transgenic mice. *Free Radical Biology & Medicine*, 51: 1831-1841. 2011.
17. Nikolaeva et al. The circadian clock modulates renal sodium handling. *J. Am. Soc. Nephrol.* 23: 1019-1026, 2012.
18. Onoe et al. Increase of 20-HETE synthase after brain ischemia in rats revealed by PET study with (11)C-labeled 20-HETE synthase-specific inhibitor. *J Cereb Blood Flow Metab* 32, 1737-1746, 2012.
19. Dong H. Metabolomic profiling of lipids for biomarker discovery. *Biochem Anal Biochem* 1, 5, 2012.
20. Fordsmann JC, Ko RW, Choi HB, Thomsen K, Witgen BM, Mathiesen C, Lønstrup M, Piilgaard H, MacVicar BA, Lauritzen M. Increased 20-HETE synthesis explains reduced cerebral blood flow but not impaired neurovascular coupling after cortical spreading depression in rat cerebral cortex. *J Neurosci* 33, 2562-2570, 2013.
21. Stobart JL, Lu L, Anderson HD, Mori H, Anderson CM. Astrocyte-induced cortical vasodilation is mediated by D-serine and endothelial nitric oxide synthase. *PNAS USA.* 110, 3149-3154, 2013.
22. Tunctan 20-HETE mimetics or inhibitors in the treatment of cancer patients with sepsis and septic shock. *International Journal of Cancer Studies & Research (IJCR)* 2:101, 2013.
23. Tunctan et al. NS-398 reverses hypotension in endotoxemic rats: Contribution of eicosanoids, NO, and peroxynitrite. *Prostaglandins & other Lipid Mediators* 104–105 93–108, 2013 .
24. Alexanian, A, Sorokin A. Targeting 20-HETE producing enzymes in cancer--rationale, pharmacology and clinical potential. *Oncotargets and Therapy* 6: 243-255, 2013.

25. Raso et al. N-Palmitoylethanolamide protects the kidney from hypertensive injury in spontaneously hypertensive rats via inhibition of oxidative stress. *Pharmacological Research* 76: 67-76, 2013.
26. Eid et al. 20-HETE and EETs in diabetic nephropathy: A novel mechanistic pathway. *PLOS One* 8: e70029, 2013
27. Ross et al. Proinflammatory high-density lipoprotein results from oxidized lipid mediators in the pathogenesis of both idiopathic and associated types of pulmonary arterial hypertension. *Pulmonary circulation* 5: 640-648, 2015