



Metro Center for High Technology Bldg.
2727 Second Ave. Suite 4113
Detroit, MI 48201
Phone: (313) 961-1606; Fax: (313)963-7130
Email: info@DetroitRandD.com
Web: www.DetroitRandD.com

Oxidative Stress ELISA (8-isoprostane) kit

(iPF2 α -III, 15-isoprostane F2t, 8-epi-prostaglandin F2 α , 8-iso-prostaglandin F2 α)

Cat # 8iso1: ELISA kit for measuring 8-isoprostane in biological / wastewater samples

Cat # 8iso1R: 8-isoprostane ELISA Kit with removable strips:


This competitive ELISA kit is for determination of 8-isoprostane levels in biological samples. A typical standard curve of the ELISA is shown in the bottom left corner.

The isoprostanes are a family of eicosanoids of non-enzymatic origin produced by the random oxidation of tissue phospholipids by oxygen radicals¹. 8-epi-prostaglandin F2a is an in vivo index of oxidative injury¹ and associated with increased coronary artery calcification (CAC)².

A recent study using the 8-isoprostane ELISA revealed that rat hepatic 8-isoprostane levels increased in high fat (HF) diet-induced non-alcoholic fatty liver disease (NAFLD). Interestingly, telmisartan, an angiotensin II receptor antagonist (hypertension drug), significantly decreased the liver 8-isoprostane levels³. Urinary excretion of 8-isoprostane increased in mice by aldosterone infusion-induced renal fibrosis. The level of 8-isoprostane excretion markedly decreased in osteopontin (involved in tumor promotion)-knock-out mice, suggesting that osteopontin is a therapeutic target for oxidative stress-induced kidney injury⁴. Levels of both renal NADPH oxidase subunit mRNA expression and urinary 8-isoprostane were increased in streptozotocin-induced diabetes mellitus (DM)⁵. 8-isoprostane has been examined as an exercise-induced muscle damage biomarker⁶ and as a sewage biomarker for assessing community health⁷. Yamamoto et al. showed that the use of AST-120 was related to improvement of oxidative stress (decrease in 8-isoprostane levels) in patients undergoing maintenance haemodialysis treatment.⁸ In utero diesel exhaust exposure of hyperlipidemic mice did not influence later life redox homeostasis⁹. A strong association between tobacco smoking and urinary 8-isoprostane levels was found in a large German study of older adult smokers^{10,11}. Soleus muscle from T2D-induced rats fed a high fat diet exhibited higher levels of 8-isoprostane in conjunction with increased mt DNA copy number and 8-OHdG¹². Hyperoxia increased 8-isoprostane in lung of wild type mice but not in soluble epoxide hydrolase knockout mice¹³.

Diluted urine or cell culture media without extraction are suitable for the ELISA. Each kit for triplicate analyses of up to 24 samples contains one 96 well plate, one tube of 8-isoprostane standard, one tube of 8-isoprostane-conjugated horseradish peroxidase (HRP), and buffers for sample and HRP dilutions, and plate washing.

Buy in Quantity and Save



References

1. MorrowJD, Hill KE, Burk RF, Nammour TM, Badr KF, Roberts LJ. A series of prostaglandin F2-like compounds are produced in vivo in humans by a non-cyclooxygenase, free radical-catalyzed mechanism. Proc Natl Acad Sci USA 87, 9383-9387, 1990.
2. Gross M, et al. Plasma F2-isoprostanes and coronary artery calcification: the CARDIA Study. Clin. Chem. 51, 125-131, 2005.
3. Sookoian et al. High fat diet-induced liver steatosis promotes an increase in liver mitochondrial biogenesis in response to hypoxia. J. Cell. Mol. Med. 15, 1329-1338, 2011.
4. Irita et al. Osteopontin deficiency protects against aldosterone-induced inflammation, oxidative stress, and interstitial fibrosis in the kidney. Am. J. Physiol. Renal Physiol. 301, F833-844 2011.
5. Nagao T, Okura T et al. Osteopontin plays a critical role in interstitial fibrosis but not glomerular sclerosis in diabetic nephropathy. Nephron Extra 2, 87-103, 2012.
6. Kanda et al. Evaluation of serum leaking enzymes and investigation into new biomarkers for exercise-induced muscle damage. EIR 20: 39-54, 2014
7. Santos et al. Could sewage epidemiology be a strategy to assess lifestyle and wellness of a large scale population? Medical Hypothesis 85: 408-411, 2015.

(continued on next page)

- /
8. Yamamoto et al. Continuous reduction of protein-bound uraemic toxins with improved oxidative stress by using the oral charcoal adsorbent AST-120 in haemodialysis patients. *Sci. Rep.* 5, 14381; doi: 10.1038/srep14381 2015
 9. Harrigan et al. In utero exposure of hyperlipidemic mice to diesel exhaust: Lack of effects on atherosclerosis in adult offspring fed a regular chow diet. *Cardiovasc Toxicol.* DOI:10.1007/s12012-017-9399, 2017.
 10. Gao et. al. Associations of self-reported smoking, cotinine levels and epigenetic smoking indicators with oxidative stress among older adults: a population-based study. *Genetic Epidemiology* 32: 443-456, 2017.
 11. Gao, X et al. Urinary isoprostane levels and occurrence of lung, colorectal, prostate, breast and overall cancer: Results from a large, population-based cohort study with 14 years of follow-up. *Free Radical Biology and Medicine* 123: 20-26. 2018.
 12. Santos, JM et al. The role of mitochondrial DNA damage at skeletal muscle oxidative stress on the development of type 2 diabetes. *Molecular and Cellular Biochemistry* doi.org/10.1007/s11010-018-3361-5. 2018.
 13. Liu, LP et al. Deletion of soluble epoxide hydrolase attenuates mice hyperoxic acute lung injury. *BMC Anesthesiology* doi.org/10.1186/s12871-018-0490-z. 2018.