

1. Schottker, B et al. Oxidatively damaged DNA/RNA and 8-isoprostane levels are associated with the development of type 2 diabetes at older age: Results from a large cohort study. *Diabetes Care*, doi: 10.2337/dc19-1379. 2019
2. Gào X, Xuan Y, Benner A, Anusruti A, Brenner H, Schöttker, B. Nitric Oxide Metabolites and Lung Cancer Incidence: A Matched Case-Control Study Nested in the ESTHER Cohort. *Oxidative Medicine and Cellular Longevity* Volume 2019, Article ID 6470950, <https://doi.org/10.1155/2019/6470950> 2
3. Kawasaki K, Kondoh E, Chigusa Y, Kawamura Y, Mogami H, Takeda S, Horie A, Baba T, Matsumura N, Mandai M, Konishi I. Metabolomic profiles of placenta in preeclampsia: Antioxidant effect of magnesium sulfate on trophoblasts in early-onset preeclampsia. *Hypertension* 73: 671-679. 2019.
4. Gào X, Zhang Y, Burwinkel B, Xuan Y, Holleccek B, Brenner H, Schöttker B. The associations of DNA methylation alterations in oxidative stress-related genes with cancer incidence and mortality outcomes: a population-based cohort study. *Clinical Epigenetics* 11: 14-22. 2019.
5. Gao, X, Gào X, Zhang Y, Holleccek B, Schöttker B, Brenner H. Oxidative stress and epigenetic mortality risk score: associations with all-cause mortality among elderly people. *Eur. J. Epidemiology* 34: 451-462. 2019.
6. Gào X, Brenner H, Holleccek B, Cuk K, Zhang Y, Anusruti A, Xuan Y, Xu Y, Schöttker B. Urinary isoprostane levels and occurrence of lung, colorectal, prostate, breast and overall cancer: Results from a large, population-based cohort study with 14 years of follow-up. *Free Radical Biology and Medicine* 123: 20-26. 2018.
7. Santos, JM et al. The role of mitochondrial DNA damage at skeletal muscle oxidative stress on the development of type 2 diabetes. *Molecular and Cellular Biochemistry* doi.org/10.1007/s11010-018-3361-5. 2018.
8. Liu, LP et al. Deletion of soluble epoxide hydrolase attenuates mice hyperoxic acute lung injury. *BMC Anesthesiology* doi.org/10.1186/s/12871-018-0490-z. 2018.
9. Harrigan et al. In utero exposure of hyperlipidemic mice to diesel exhaust: Lack of effects on atherosclerosis in adult offspring fed a regular chow diet. *Cardiovasc Toxicol*. DOI:10.1007/s12012-017-9399, 2017.
10. Gao et. al. Associations of self-reported smoking, cotinine levels and epigenetic smoking indicators with oxidative stress among older adults: a population-based study. *Genetic Epidemiology* 32: 443- 456, 2017.
11. Yamamoto et al. Continuous reduction of protein-bound uraemic toxins with improved oxidative stress by using the oral charcoal adsorbent AST-120 in haemodialysis patients. *Sci. Rep.* 5, 14381; doi: 10.1038/srep14381 2015.
12. Santos et. al. Could sewage epidemiology be a strategy to assess lifestyle and wellness of a large scale population? *Medical Hypothesis* 85: 408-411, 2015.

13. Kanda et al. Evaluation of serum leaking enzymes and investigation into new biomarkers for exercise-induced muscle damage. *EIR* 20: 39-54, 2014
14. Nagao T, Okura T et al. Osteopontin plays a critical role in interstitial fibrosis but not glomerular sclerosis in diabetic nephropathy. *Nephron Extra* 2, 87-103, 2012.
15. Irita et al. Osteopontin deficiency protects against aldosterone-induced inflammation, oxidative stress, and interstitial fibrosis in the kidney. *Am. J. Physiol. Renal Physiol.* 301, F833-844, 2011.
16. Sookoian et al. High fat diet-induced liver steatosis promotes an increase in liver mitochondrial biogenesis in response to hypoxia. *J. Cell. Mol. Med.* 15, 1329-1338, 2011.