

## Rabbit Anti-Transferrin (TF)


(100 µg/vial, Lyophilized)

**Catalog Number: BPAB3661**


FOR RESEARCH USE ONLY

**Detroit R&D, Inc.**


<b>Product Name</b>	Anti-Transferrin/TF Antibody
<b>Reactive Species</b>	Human, Mouse, Rat
<b>Description</b>	Rabbit IgG polyclonal antibody for Serotransferrin(TF) detection. Tested with WB, IHC-P in Human;Mouse;Rat.
<b>Application</b>	IHC, WB
<b>Clonality</b>	Polyclonal
<b>Formulation</b>	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg NaN <sub>3</sub> .
<b>Storage Instructions</b>	At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.
<b>Host</b>	Rabbit
<b>Uniprot ID</b>	P02787
<b>Immunogen</b>	A synthetic peptide corresponding to a sequence at the N-terminus of human Transferrin.
<b>Cross Reactivity</b>	No cross reactivity with other proteins
<b>Form</b>	Lyophilized
<b>Concentration</b>	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
<b>Purification</b>	Immunogen affinity purified.
<b>Suggested Dilutions</b>	Immunohistochemistry(Paraffin-embedded Section), 0.5-1µg/ml, Human, Mouse, Rat, By Heat Western blot, 0.1-0.5µg/ml, Human, Rat




Anti- Transferrin antibody Western blotting  
All lanes: Anti Transferrin (at 0.5ug/ml)  
Lane 1: Rat Thymus Tissue Lysate at 50ug  
Lane 2: Human Placenta Tissue Lysate at 50ug  
Predicted bind size: 77KD  
Observed bind size: 77KD



Anti-Transferrin antibody  
IHC(P): Mouse Liver Tissue



Anti-Transferrin antibody  
IHC(P): Rat Liver Tissue



Anti-Transferrin antibody  
IHC(P): Human Intestinal Cancer Tissue