



Metro Center for High Technology Bldg.
2727 Second Ave. Suite 4113
Detroit, MI 48201
Phone: (313) 961-1606; Fax: (313)963-7130
Email: info@DetroitRandD.com

Hypertension ELISA (11,12-DHET) kit

Cat # DH 4: ELISA kit for measuring 11,12-DHET in biological samples

This competitive ELISA kit is for determination of 11,12-DHET levels in biological samples¹. Anti-11,12-DHET does not cross-react with 5(S)15(S)DiHETE and arachidonic acid (AA) and minimally cross-reacts with 8,9- and 14,15-DHETs. The 11,12-DHET is a metabolite of soluble epoxide hydrolase (sEH)-mediated metabolism of the 11,12-EET, which is generated by AA epoxygenase activity of cytochromes P450. The 11,12-DHET plays a critical role in vasorelaxation mediated by endothelial cell large-conductance Ca^{2+} -activated K^{+} (BK_{Ca}) channels^{2,3}. Whereas 8,9- and 14,15-DHETs dilated isolated human coronary arterioles (HCAs) less than the corresponding EETs, 11,12-DHET dilated them similarly to the 11,12-EET⁴. 11,12-DHET ELISA was used to quantify eicosapentaenoic acid (EPA)-dependent increase of 11,12-EET levels (via PPAR γ) in endothelial cells⁵. High glucose reduced 11,12-EET production in porcine coronary artery endothelial cells, which impaired EET/endothelium-derived hyperpolarizing factor (EDHF)-mediated vasodilation in coronary arterioles⁶. Recently, 11,12-EETs were found to protect against angiotensin II-induced abdominal aortic aneurysm in mice⁷ and increased after indapamide treatment, which lowered blood pressure⁸.

Each kit for triplicate analyses of up to 24 samples contains one 96 well plate, one tube of 11,12-DHET standard, one tube of 11,12-DHET-conjugated horseradish peroxidase (HRP), and buffers for sample and HRP dilutions, and plate washing.

Cat # DH 14: 2X ELISA kits for measuring 11,12-DHET in biological samples

Cat # DH 24: 5X ELISA kits for measuring 11,12-DHET in biological samples


Cat # DH 104: 10X ELISA kits for measuring 11,12-DHET in biological samples

Two Other Hypertension ELISA kits:

- 20 HETE Hypertension ELISA kit
- 14,15-DHET Hypertension ELISA kit

Oxidative Stress ELISA Kit:

- 8-iso PGF_{2 α} (isoprostane) ELISA kit



1. Kim et al. Three US Patents: 14,15 and 11,12-DHET dependent hypertension 6,440,682, 6,534,282 and 7,695,927 issued on 8/27/2002, 3/18/2003 & 4/13/2010 respectively. (<http://patft.uspto.gov/netahtml/srchnum.htm>)
2. X. Fang, T.L. Kaduce, N.L. Weintraub, M. VanRollins, A.A. Spector. Functional implications of a newly characterized pathway of 11,12-epoxyeicosatrienoic acid metabolism in arterial smooth muscle. *Circ. Res.* 79, 784-804, 1996.
3. C.L. Oltman, N.L. Weintraub, M. VanRollins, K.C. Dellsperger. Epoxyeicosatrienoic acid and dihydroxyeicosatrienoic acids are potent vasodilators in the canine coronary microcirculation. *Circ. Res.* 83, 932-939, 1998.
4. B.T. Larsen, H. Miura, O.A. Hatoum, W.B. Campbell, B.D. Hammock, D.C. Zeldin, J.R. Falck, and D.D. Gutterman. Epoxyeicosatrienoic acid and dihydroxy-eicosatrienoic acids dilate human coronary arterioles via BKCa channels: implications for soluble epoxide hydrolase inhibition. *Am. J. Physiol. Heart Circ. Physiol.* 290, 491-499, 2006.
5. D. Wang, T. Hirase, T. Nitto, M. Soma, K. Node. Eicosapentaenoic acid increases cytochrome P450 2J2 gene expression and epoxyeicosatrienoic acid production via peroxisome proliferator-activated receptor γ in endothelial cells. *J. Cardiol.* 54, 368-374, 2009.
6. Yang, Kuo et al., High glucose impairs EDHF-mediated dilation of coronary arterioles via reduced cytochrome P450 activity. *Microvasc. Res.* 82, 356-363, 2011.
7. Cai, Wang et al. CYP2J2 overexpression increases EETs and protects against angiotensin II-induced abdominal aortic aneurysm in mice. *J lipid Res* 54, 1448-1456, 2013.
8. Ma F, Lin F, Chen C, Cheng J, Zeldin DC, Wang Y, Wang DW. Indapamide lowers blood pressure by increasing production of epoxyeicosatrienoic acids in the kidney. *Mol Pharmacol* 84,286-295, 2013