



**CAT # GDF-15**

**Polyclonal Goat Anti-Human GDF-15 (NAG-1, MIC-1) Antibody**

(50  $\mu$ L)

**Preparation:** Polyclonal antibody for human GDF-15/PTGF $\beta$  was produced by immunization of a rabbit with a human NAG-1 peptide antigen.

**Specificity:** The antibody recognizes pro- and secreted- forms of GDF-15/PTGF $\beta$  in human.

**Contents:** 50  $\mu$ L per vial with 0.7% BSA and 0.05% azide.

**Storage:** Avoid repeated freezing and thawing by making aliquots of the antibody solution and storing at  $-20^{\circ}\text{C}$ .

**Western blots:** 5  $\mu$ L per 10 mL of TBS buffer (1:2,000 dilution) containing 5% non-fat milk for overnight incubation is necessary for ECL method. We recommend a 1:5,000 dilution of HRP conjugated anti-rabbit secondary antibody in TBS containing 5% non-fat milk for 1 hr incubation.

Metro Center for High Tech Bldg.  
2727 Second Ave. Suite #4113  
Detroit, MI 48201  
Phone: (313) 961-1606  
Fax: (313) 963-7130  
email: [info@DetroitRandD.com](mailto:info@DetroitRandD.com)  
Web: DetroitRandD.com

Revised 26 October 2016